

**AMENDMENTS TO THE CLAIMS**

Please amend the Claim Form and Claim as follows. Insertions are shown underlined while deletions are ~~struck through~~.

1 (canceled)

~~2~~ <sup>2</sup> (previously presented): The method according to Claim 10, wherein said nuclease is a nuclease contained in the yeast RNA-containing composition.

~~3~~ <sup>3</sup> (previously presented): The method according to Claim 10, wherein the yeast RNA-containing composition is obtained from yeast selected from the group consisting of *Saccharomyces cerevisiae* and *Candida utilis*.

4 (currently amended): The method according to Claim 15, wherein the decomposition step is conducted by digesting the yeast RNA-containing composition with ribonuclease A and/or trypsin~~added to a solution containing the yeast RNA-containing composition, at a pH value of 3-10 and at a temperature of 10-70°C.~~ <sup>14</sup>

~~10~~ <sup>10</sup> (currently amended): The method according to Claim ~~15~~ <sup>14</sup>, wherein the decomposition step is conducted by hydrolyzing at 20-100°C the yeast RNA-containing composition with sodium hydrate~~alkali added to a solution containing the yeast RNA-containing composition at a normality of 0.1-5N.~~

~~4~~ <sup>4</sup> (previously presented): The method according to Claim 10, wherein the yeast RNA-containing composition is an extract obtained by physically crushing yeast using a high-pressure homogenizer and an ultrasonic disintegrator.

~~5~~ <sup>5</sup> (previously presented): The method according to Claim 10, wherein the yeast RNA-containing composition is an extract obtained from yeast using hot water at a pH value of 4-8 and at a temperature of 90-100°C, wherein sodium chloride is added to a yeast suspension with a yeast concentration of 5-25% to make a salt concentration of 1-10%.

~~6~~ <sup>6</sup> (previously presented): The method according to Claim 10, wherein the yeast RNA-containing composition is an extract obtained by autolyzing yeast.

9 (canceled)

~~10~~ <sup>10</sup> (currently amended): A method of obtaining polyamines, comprising the steps of:  
providing a yeast RNA-containing composition;  
subjecting said yeast RNA-containing composition to a decomposition step,  
comprising nuclease digestion or alkali hydrolysis, ~~effor~~ for increasing the yield of

polyamines recovered in a subsequent recovery step by approximately 2-3.2 times the yield of polyamines recovered in the subsequent recovery step without this decomposition step, under conditions where the yield with this decomposition step when continuing for approximately 15-18 hours is approximately 2-3.2 times the yield without this decomposition step, wherein said yeast RNA-containing composition is treated in solution with nuclease added in an effective concentration of approximately 1-2 mg/ml, at approximately 25-37°C, and at a pH of approximately 6-8, or said yeast RNA-containing composition is dissolved in a 0.3 N alkali solution at 37°C; and

recovering the approximately 2-3.2 times greater yield of polyamines from the decomposed yeast RNA-containing composition produced.

7 ~~11~~ (currently amended): The method according to Claim 10, wherein the nuclease is selected from the group consisting of ~~deoxyribonuclease I~~, nuclease P1, nuclease S1, phosphodiesterase I, ribonuclease A, ribonuclease B, ribonuclease T<sub>1</sub>, ribonuclease T<sub>2</sub>, and ribonuclease U<sub>2</sub>.

8 ~~12~~ (previously presented): The method according to Claim 10, wherein the alkali is sodium hydrate or potassium hydroxide.

13 (canceled)

9 ~~14~~ (currently amended): A method of obtaining polyamines, comprising ~~the steps of:~~  
a step for providing a yeast RNA-containing composition;  
a step for subjecting said yeast RNA-containing composition to a decomposition step, comprising nuclease digestion or alkali hydrolysis, ~~effor~~ increasing the yield of polyamines recovered in a subsequent recovery step by approximately 2-3.2 times the yield of polyamines recovered in the subsequent recovery step without this decomposition step, under conditions where said yeast RNA-containing composition is treated in solution with nuclease added in an effective concentration of approximately 1-2 mg/ml, at approximately 25-37°C, and at a pH of approximately 6-8, or said yeast RNA-containing composition is dissolved in a 0.3 N alkali solution at 37°C; and

a step for recovering the approximately 2-3.2 times greater yield of polyamines from the decomposed yeast RNA-containing composition produced.

15 (currently amended): A method of obtaining polyamines, comprising the steps of:  
providing a yeast RNA-containing composition;

Appl. No. : 09/514,999  
Filed : February 29, 2000

decomposing said yeast RNA-containing composition by nuclease digestion or alkali hydrolysis to ~~separate~~yield polyamines from high-molecular weight substances in the yeast RNA-containing composition ~~to a degree achieved in an amount obtained by treating when the yeast RNA-containing composition is treated~~ for about 15-18 hours in solution with nuclease added in an effective ~~concentration of about 1-2 mg/ml~~, at about 25-37°C, and at a pH of about 6-8, or in an about 0.3 N alkali solution at about 37°C; and recovering polyamines from the decomposed yeast RNA-containing composition produced.